



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

262. Proposed by NELSON L. RORAY, Utica, New York.

In a regular pentagon, show that $\frac{\text{diagonal}}{\text{side}} = \frac{2 \text{ apothem}}{\text{radius}} = \frac{5 + \sqrt{5}}{\sqrt{5}}$.

CALCULUS.

199. Proposed by F. P. MATZ, Sc. D., Ph. D., Reading, Pa.

If the perimeter of an ellipse varies uniformly at the rate of $\frac{1}{4}$ inch per unit of time, at what rate is the eccentricity varying the instant the perimeter becomes 60 inches and the major axis 25 inches?

200. Proposed by R. D. CARMICHAEL, Hartselle, Alabama.

Find the equation of a curve so that the area bounded by the curve, the axis of x , and any ordinate y , is equal to $y-x$, x being the *corresponding* abscissa.

201. Proposed by F. P. MATZ, Sc. D., Ph. D., Reading, Pa.

Solve $\iint \frac{dy/dx}{1+w^2} dw = 0$.

MECHANICS.

180. Proposed by EDWIN L. RICH, Lehigh University, South Bethlehem, Pa.

If a body is projected into the air, and the resistance of the air varies as the square of the velocity; required the equation of the curve. [From De Volson Wood's *Analytical Mechanics*, problem 10, p. 179.]

181. Proposed by F. ANDEREGG, Professor of Mathematics, Oberlin College, Oberlin, Ohio.

A triangle AOB , of which the sides, OA , AB , and the angle at O are a , b , and α , revolves uniformly about O , so that OA makes the angle nt with the axis of x , and carries a circle of which AB is the diameter. Prove that a point moving in the circumference of the carried circle with twice the angular velocity of the triangle will describe an ellipse whose axes are

$$\sqrt{(a^2 + b^2 + 2ab \cos \alpha)} \pm \sqrt{(a^2 + b^2 - 2ab \cos \alpha)}.$$

182. Proposed by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

I have a tank, the lower part of which is a hemisphere 22 feet in diameter. The rest is a cylinder 22 feet in diameter, and altitude 28 feet. This tank is connected with the earth by a vertical stand-pipe 10 inches in diameter, 130 feet long, extending 2 feet into the tank. The tank is filled by a $2\frac{1}{2}$ inch pipe 65 feet long, having one right-angled elbow delivering the water into the bottom of the stand-pipe from a steam pump under 96 pounds gauge pressure. How long will it take to fill the pipe?